

MATHEMATICS

PAPER—I

Time Allowed : Three Hours

Maximum Marks : 200

QUESTION PAPER SPECIFIC INSTRUCTIONS

**Please read each of the following instructions carefully
before attempting questions**

There are EIGHT questions in all, out of which FIVE are to be attempted.

Question Nos. 1 and 5 are compulsory. Out of the remaining SIX questions, THREE are to be attempted selecting at least ONE question from each of the two Sections A and B.

The number of marks carried by a question/part is indicated against it.

Unless otherwise mentioned, symbols and notations have their usual standard meanings.

Assume suitable data, if necessary, and indicate the same clearly.

Attempts of questions shall be counted in sequential order. Unless struck off, attempt of a question shall be counted even if attempted partly. Any page or portion of the page left blank in the Question-cum-Answer Booklet must be clearly struck off.

Answers must be written in ENGLISH only.

SECTION—A

1. (a) Let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a linear operator on \mathbb{R}^3 defined by $T(x, y, z) = (2y + z, x - 4y, 3x)$. Find the matrix of T in the basis $\{(1, 1, 1), (1, 1, 0), (1, 0, 0)\}$. 8
- (b) The eigenvalues of a real symmetric matrix A are $-1, 1$ and -2 . The corresponding eigenvectors are $\frac{1}{\sqrt{2}}(-1 \ 1 \ 0)^T$, $(0 \ 0 \ 1)^T$ and $\frac{1}{\sqrt{2}}(-1 \ -1 \ 0)^T$ respectively. Find the matrix A^4 . 8
- (c) Find the volume lying inside the cylinder $x^2 + y^2 - 2x = 0$ and outside the paraboloid $x^2 + y^2 = 2z$, while bounded by xy -plane. 8
- (d) Justify by using Rolle's theorem or mean value theorem that there is no number k for which the equation $x^3 - 3x + k = 0$ has two distinct solutions in the interval $[-1, 1]$. 8
- (e) If the coordinates of the points A and B are respectively $(b\cos\alpha, b\sin\alpha)$ and $(a\cos\beta, a\sin\beta)$ and if the line joining A and B is produced to the point $M(x, y)$ so that $AM : MB = b : a$, then show that $x\cos\frac{\alpha+\beta}{2} + y\sin\frac{\alpha+\beta}{2} = 0$. 8

2. (a) Determine the extreme values of the function $f(x, y) = 3x^2 - 6x + 2y^2 - 4y$ in the region $\{(x, y) \in \mathbb{R}^2 : 3x^2 + 2y^2 \leq 20\}$. 10
- (b) Consider the singular matrix

$$A = \begin{bmatrix} -1 & 3 & -1 & 1 \\ -3 & 5 & 1 & -1 \\ 10 & -10 & -10 & 14 \\ 4 & -4 & -4 & 8 \end{bmatrix}$$

Given that one eigenvalue of A is 4 and one eigenvector that does not correspond to this eigenvalue 4 is $(1 \ 1 \ 0 \ 0)^T$. Find all the eigenvalues of A other than 4 and hence also find the real numbers p, q, r that satisfy the matrix equation $A^4 + pA^3 + qA^2 + rA = 0$. 15

- (c) A line makes angles $\alpha, \beta, \gamma, \delta$ with the four diagonals of a cube. Show that

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma + \cos^2 \delta = \frac{4}{3} \quad 15$$

3. (a) Consider the vectors $x_1 = (1, 2, 1, -1)$, $x_2 = (2, 4, 1, 1)$, $x_3 = (-1, -2, 0, -2)$ and $x_4 = (3, 6, 2, 0)$ in \mathbb{R}^4 . Justify that the linear span of the set $\{x_1, x_2, x_3, x_4\}$ is a subspace of \mathbb{R}^4 , defined as

$$\{(\xi_1, \xi_2, \xi_3, \xi_4) \in \mathbb{R}^4 : 2\xi_1 - \xi_2 = 0, 2\xi_1 - 3\xi_3 - \xi_4 = 0\}$$

Can this subspace be written as $\{(\alpha, 2\alpha, \beta, 2\alpha - 3\beta) : \alpha, \beta \in \mathbb{R}\}$? What is the dimension of this subspace?

15

- (b) The dimensions of a rectangular box are linear functions of time— $l(t)$, $w(t)$ and $h(t)$. If the length and width are increasing at the rate 2 cm/sec and the height is decreasing at the rate 3 cm/sec, find the rates at which the volume V and the surface area S are changing with respect to time. If $l(0) = 10$, $w(0) = 8$ and $h(0) = 20$, is V increasing or decreasing, when $t = 5$ sec? What about S , when $t = 5$ sec?

10

- (c) Show that the shortest distance between the straight lines

$$\frac{x-3}{3} = \frac{y-8}{-1} = \frac{z-3}{1} \quad \text{and} \quad \frac{x+3}{-3} = \frac{y+7}{2} = \frac{z-6}{4}$$

is $3\sqrt{30}$. Find also the equation of the line of shortest distance.

15

4. (a) Using elementary row operations, reduce the matrix

$$A = \begin{bmatrix} 2 & 1 & 3 & 0 \\ 3 & 0 & 2 & 5 \\ 1 & 1 & 1 & 1 \\ 2 & 1 & 1 & 3 \end{bmatrix}$$

to reduced echelon form and find the inverse of A and hence solve the system of linear equations $AX = b$, where $X = (x, y, z, u)^T$ and $b = (2, 1, 0, 4)^T$.

15

- (b) Find the centroid of the solid generated by revolving the upper half of the cardioid $r = a(1 + \cos\theta)$ bounded by the line $\theta = 0$ about the initial line. Take the density of the solid as uniform.

10

- (c) A variable plane is parallel to the plane $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 0$ and meets the axes at the points A , B and C . Prove that the circle ABC lies on the cone

$$yz \left(\frac{b}{c} + \frac{c}{b} \right) + zx \left(\frac{c}{a} + \frac{a}{c} \right) + xy \left(\frac{a}{b} + \frac{b}{a} \right) = 0$$

15

SECTION—B

5. (a) Solve the differential equation $(D^2 + 1)y = x^2 \sin 2x$; $D \equiv \frac{d}{dx}$. 8

(b) Solve the differential equation $(px - y)(py + x) = h^2 p$, where $p = y'$. 8

(c) A 2 metres rod has a weight of 2 N and has its centre of gravity at 120 cm from one end. At 20 cm, 100 cm and 160 cm from the same end are hung loads of 3 N, 7 N and 10 N respectively. Find the point at which the rod must be supported if it is to remain horizontal. 8

(d) Let $\bar{r} = \bar{r}(s)$ represent a space curve. Find $\frac{d^3 \bar{r}}{ds^3}$ in terms of \bar{T} , \bar{N} and \bar{B} , where \bar{T} , \bar{N} and \bar{B} represent tangent, principal normal and binormal respectively. Compute $\frac{d\bar{r}}{ds} \cdot \left(\frac{d^2 \bar{r}}{ds^2} \times \frac{d^3 \bar{r}}{ds^3} \right)$ in terms of radius of curvature and the torsion. 8

(e) Evaluate $\int_{(0,0)}^{(2,1)} (10x^4 - 2xy^3)dx - 3x^2y^2dy$ along the path $x^4 - 6xy^3 = 4y^2$. 8

6. (a) Solve by the method of variation of parameters the differential equation

$$x''(t) - \frac{2x(t)}{t^2} = t, \quad \text{where } 0 < t < \infty \quad 15$$

(b) Find the law of force for the orbit $r^2 = a^2 \cos 2\theta$ (the pole being the centre of the force). 15

(c) Verify Stokes' theorem for $\bar{V} = (2x - y)\hat{i} - yz^2\hat{j} - y^2z\hat{k}$, where S is the upper half surface of the sphere $x^2 + y^2 + z^2 = 1$ and C is its boundary. 10

7. (a) Find the general solution of the differential equation

$$\ddot{x} + 4x = \sin^2 2t$$

Hence find the particular solution satisfying the conditions

$$x\left(\frac{\pi}{8}\right) = 0 \quad \text{and} \quad \dot{x}\left(\frac{\pi}{8}\right) = 0 \quad 15$$

- (b) A vessel is in the shape of a hollow hemisphere surmounted by a cone held with the axis vertical and vertex uppermost. If it is filled with a liquid so as to submerge half the axis of the cone in the liquid and height of the cone be double the radius (r) of its base, find the resultant downward thrust of the liquid on the vessel in terms of the radius of the hemisphere and density (ρ) of the liquid. 15
- (c) Derive the Frenet-Serret formulae. Verify the same for the space curve $x = 3 \cos t$, $y = 3 \sin t$, $z = 4t$. 10
8. (a) Find the general solution of the differential equation

$$(x-2)y'' - (4x-7)y' + (4x-6)y = 0$$
 10
- (b) A shot projected with a velocity u can just reach a certain point on the horizontal plane through the point of projection. So in order to hit a mark h metres above the ground at the same point, if the shot is projected at the same elevation, find increase in the velocity of projection. 15
- (c) Derive $\nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ in spherical coordinates and compute

$$\nabla^2 \left(\frac{x}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \right)$$
 in spherical coordinates. 15

★ ★ ★

MATHEMATICS
Paper – IITime Allowed : **Three Hours**Maximum Marks : **200****Question Paper Specific Instructions**

Please read each of the following instructions carefully before attempting questions :

*There are **EIGHT** questions in all, out of which **FIVE** are to be attempted.*

*Questions no. **1** and **5** are **compulsory**. Out of the remaining **SIX** questions, **THREE** are to be attempted selecting at least **ONE** question from each of the two Sections A and B.*

Attempts of questions shall be counted in sequential order. Unless struck off, attempt of a question shall be counted even if attempted partly. Any page or portion of the page left blank in the Question-cum-Answer Booklet must be clearly struck off.

All questions carry equal marks. The number of marks carried by a question/part is indicated against it.

*Answers must be written in **ENGLISH** only.*

Unless otherwise mentioned, symbols and notations have their usual standard meanings.

Assume suitable data, if necessary, and indicate the same clearly.

SECTION A

- Q1.** (a) Let R be an integral domain. Then prove that $\text{ch } R$ (characteristic of R) is 0 or a prime. 8
- (b) Show that the function $f(x) = \sin\left(\frac{1}{x}\right)$ is continuous and bounded in $(0, 2\pi)$, but it is not uniformly continuous in $(0, 2\pi)$. 8
- (c) Test the Riemann integrability of the function f defined by
- $$f(x) = \begin{cases} 0 & \text{when } x \text{ is rational} \\ 1 & \text{when } x \text{ is irrational} \end{cases}$$
- on the interval $[0, 1]$. 8
- (d) Using Cauchy's Integral formula, evaluate the integral $\oint_c \frac{dz}{(z^2 + 4)^2}$ where $c : |z - i| = 2$. 8
- (e) A firm manufactures two products A and B on which the profits earned per unit are ₹ 3 and ₹ 4 respectively. Each product is processed on two machines M1 and M2. Product A requires one minute of processing time on M1 and two minutes on M2, while B requires one minute on M1 and one minute on M2. Machine M1 is available for not more than 7 hours 30 minutes, while machine M2 is available for 10 hours during any working day. Find the number of units of products A and B to be manufactured to get maximum profit, using graphical method. 8
- Q2.** (a) Let I and J be ideals in a ring R . Then prove that the quotient ring $(I + J)/J$ is isomorphic to the quotient ring $I/(I \cap J)$. 10
- (b) Show that the integral $\int_0^{\pi/2} \log \sin x \, dx$ is convergent and hence evaluate it. 15
- (c) If $f(z)$ is analytic in a domain D and $|f(z)|$ is a non-zero constant in D , then show that $f(z)$ is constant in D . 15

- Q3.** (a) If in the group G , $a^5 = e$, $aba^{-1} = b^2$ for some $a, b \in G$, find the order of b . 10
- (b) Show that the sequence $\{\tan^{-1} nx\}$, $x \geq 0$ is uniformly convergent on any interval $[a, b]$, $a > 0$ but is only pointwise convergent on $[0, b]$. 15
- (c) Use simplex method to solve the following problem : 15
- Maximize $z = 2x_1 + 5x_2$
- subject to $x_1 + 4x_2 \leq 24$
- $3x_1 + x_2 \leq 21$
- $x_1 + x_2 \leq 9$
- $x_1, x_2 \geq 0$

- Q4.** (a) Show that the smallest subgroup V of A_4 containing $(1, 2)(3, 4)$, $(1, 3)(2, 4)$ and $(1, 4)(2, 3)$ is isomorphic to the Klein 4-group. 10
- (b) Classify the singular point $z = 0$ of the function $f(z) = \frac{e^z}{z + \sin z}$ and obtain the principal part of the Laurent series expansion of $f(z)$. 15
- (c) A salesman wants to visit cities $C1, C2, C3$ and $C4$. He does not want to visit any city twice before completing the tour of all the cities and wishes to return to his home city, the starting station. Cost of going from one city to another in rupees is given below in the table. Find the least cost route. 15

		To City			
		C1	C2	C3	C4
From City	C1	0	30	80	50
	C2	40	0	140	30
	C3	40	50	0	20
	C4	70	80	130	0

SECTION B

- Q5.** (a) Find the solution of the equation :

$$\frac{\partial^2 z}{\partial x^2} - \frac{\partial^2 z}{\partial y^2} = x - y.$$

8

- (b) The following table gives the values of $y = f(x)$ for certain equidistant values of x . Find the value of $f(x)$ when $x = 0.612$ using Newton's forward difference interpolation formula.

8

$x :$	0.61	0.62	0.63	0.64	0.65
$y = f(x) :$	1.840431	1.858928	1.877610	1.896481	1.915541

- (c) Following values of x_i and the corresponding values of y_i are given. Find

$$\int_0^3 y \, dx \text{ using Simpson's one-third rule.}$$

8

$x_i :$	0.0	0.5	1.0	1.5	2.0	2.5	3.0
$y_i :$	0.0	0.75	1.0	0.75	0.0	-1.25	-3.0

- (d) Consider the flow field given by $\psi = a(x^2 - y^2)$, 'a' being a constant. Show that the flow is irrotational. Determine the velocity potential for this flow and show that the streamlines and equivelocity potential curves are orthogonal.

8

- (e) Find a complete integral of the equation by Charpit's method $p^2x + q^2y = z$. Here $p = \frac{\partial z}{\partial x}$, $q = \frac{\partial z}{\partial y}$.

8

- Q6.** (a) Test the integrability of the equation

$$z(z + y^2) \, dx + z(z + x^2) \, dy - xy(x + y) \, dz = 0.$$

If integrable, then find its solution.

15

- (b) Solve the following system of equations by Gauss-Jordan elimination method :

10

$$x_1 + x_2 + x_3 = 3$$

$$2x_1 + 3x_2 + x_3 = 6$$

$$x_1 - x_2 - x_3 = -3$$

- (c) For a dynamical system

$$T = \frac{1}{2} \{ (1 + 2k) \dot{\theta}^2 + 2\dot{\theta}\dot{\phi} + \dot{\phi}^2 \},$$

$$V = \frac{n^2}{2} \{ (1 + k) \theta^2 + \phi^2 \},$$

where θ, ϕ are coordinates and n, k are positive constants, write down the Lagrange's equations of motion and deduce that

$$(\ddot{\theta} - \ddot{\phi}) + n^2 \left(\frac{1+k}{k} \right) (\theta - \phi) = 0.$$

Further show that if $\theta = \phi, \dot{\theta} = \dot{\phi}$ at $t = 0$, then $\theta = \phi$ for all t .

15

- Q7.** (a) Given $\frac{dy}{dx} = x^2 + y^2, y(0) = 1$. Find $y(0.1)$ and $y(0.2)$ by fourth order Runge-Kutta method.

15

- (b) Consider a mass-spring system consisting of a mass m and a linear spring of stiffness k hanging from a fixed point. Find the equation of motion using the Hamiltonian method, assuming that the displacement x is measured from the unstretched position of the string.

10

- (c) Find the equations of the system of curves on the cylinder $2y = x^2$ orthogonal to its intersections with the hyperboloids of the one-parameter system $xy = z + c$.

15

- Q8.** (a) Consider that the region $0 \leq z \leq h$ between the planes $z = 0$ and $z = h$ is filled with viscous incompressible fluid. The plane $z = 0$ is held at rest and the plane $z = h$ moves with constant velocity $V \hat{j}$. When conditions are steady, assuming there is no slip between the fluid and either boundary, and neglecting body forces, show that the velocity profile between the plates is parabolic. Find the tangential stress at any point $P(x, y, z)$ of the fluid and determine the drag per unit area on both the planes.

15

- (b) State the Newton–Raphson iteration formula to compute a root of an equation $f(x) = 0$ and hence write a program in BASIC to compute a root of the equation

$$\cos x - xe^x = 0$$

lying between 0 and 1. Use DEF function to define $f(x)$ and $f'(x)$.

10

- (c) Use Gauss quadrature formula of point six to evaluate $\int_0^1 \frac{dx}{1+x^2}$ given

$$x_1 = -0.23861919, \quad w_1 = 0.46791393$$

$$x_2 = -0.66120939, \quad w_2 = 0.36076157$$

$$x_3 = -0.93246951, \quad w_3 = 0.17132449$$

$$x_4 = -x_1, \quad x_5 = -x_2, \quad x_6 = -x_3, \quad w_4 = w_1, \quad w_5 = w_2 \quad \text{and} \quad w_6 = w_3.$$

15